



ORAL PRESENTATION

Open Access

O20 - Human rhinovirus replication-dependent induction of micro-RNAs in human bronchial epithelial cells

Spyridon Megremis^{1*}, Styliani Taka¹, Anastasis Oulas², Georgios Kotoulas², Ioannis Iliopoulos³, Nikolaos G Papadopoulos¹

From 3rd Pediatric Allergy and Asthma Meeting (PAAM)
Athens, Greece. 17-19 October 2013

Background

Micro-RNAs (miRNAs) are a class of small non-coding RNA molecules that function through post transcriptional regulation of gene expression by a process termed RNA interference (RNAi). RNAi-mediated targeting of viral RNAs is recognized as an antiviral defense mechanism. The epigenetic effect of miRNAs can either be direct, by interfering with virus genome, or indirect, through down-regulation of type I IFN genes. The aim of this study is to identify HRV-A1B specific miRNAs in human bronchial cell line.

Method

In silico prediction of potential HRV-A1B specific human mature miRNAs was performed using two different prediction tools, miRBase and RNAhybrid. Human bronchial epithelial cells (BEAS-2B) were infected with HRV-A1B (1 MOI) along with UV inactivated HRV1B (1 MOI), zymosan (TLR4 stimulator) and Poly I:C (TLR3 stimulator). RNA was isolated at different time points and the kinetics of 8 miRNAs were evaluated. The expression of miRNAs was measured by miRNA specific RT-QPCR. The results were calculated according to the $2^{-\Delta\Delta CT}$ method (FI). Statistical analysis was performed using Student's t test.

Results

Sixty two miRNAs were predicted to bind to the HRV-A1B positive strand. Eight miRNAs were selected according to their binding properties. We found replication dependent

HRV-A1B specific induction in hs-miR-a (50 FI) and miR-b (24 FI) at 7 hours after HRV1B infection.

Conclusion

To our knowledge, this is the first study to demonstrate replication dependent induction of HRV-A1B specific human miRNAs in human bronchial epithelial cell line. The expression levels of hs-miR-a and hs-miR-b were HRV replication-dependent. Further experiments are needed in order to define the potential antiviral activity of the above miRNAs.

Authors' details

¹Allergy Department, ²nd Pediatric Clinic, University of Athens, Athens, Greece. ²Institute of Marine Biology, Biotechnology and Aquaculture - HCMR, Heraklion, Crete, Greece. ³Division of Medical Sciences, University of Crete, Heraklion, Crete, Greece.

Published: 28 February 2014

doi:10.1186/2045-7022-4-S1-O20

Cite this article as: Megremis et al.: O20 - Human rhinovirus replication-dependent induction of micro-RNAs in human bronchial epithelial cells. *Clinical and Translational Allergy* 2014 **4**(Suppl 1):O20.

¹Allergy Department, ²nd Pediatric Clinic, University of Athens, Athens, Greece

Full list of author information is available at the end of the article