

POSTER PRESENTATION

Open Access

A case of adult eosinophilic oesophagitis

Isabel Skypala^{1*}, Stephen Till²

From Food Allergy and Anaphylaxis Meeting 2011 Venice, Italy. 17-19 February 2011

Background

Adults with Eosinophilic Oesophagitis (EO) often present with a history of dysphagia and food impaction. Many are also atopic and sensitisation to foods is common; 81% of adults with EO in one study were sensitised to one or more food or inhalant allergen, and 50% had positive skin prick tests to one or more foods. Exclusion of foods has been reported to improve clinical symptoms; 94% of adults improved on a 6 food elimination diet, and in children, the use of an elemental formula has been proven to be effective.

Method

We describe an adult male patient aged 30 years with a diagnosis of EO. Skin prick testing with fresh foods and specific IgE blood tests (Phadia ImmunoCAP) revealed sensitisation to a number of foods. Following an acute episode of dysphagia requiring dilatation, the patient commenced a complete exclusion diet. Nutritional support was maintained solely through the consumption of a nutritionally complete liquid elemental diet (Elemental 028™). After six weeks, one new food was introduced every 2-3 days.

Results

The patient was sensitised to birch and grass pollen and a wide variety of foods including potatoes, carrots, broccoli, lettuce, spinach, beer, wine, grapes, apple juice, mango, peanut, mustard, marmite, rice, wheat, barley, yeast and the peach lipid transfer protein allergen Pru p 3. Following the 6-week elemental diet, to date the following foods have been successfully re-introduced into the diet; rice, fish, onions, eggs, olives, lamb, corn, barley, milk, vodka, red wine, champagne, barley, peas, pak choi, pork, turkey. Soy and grapes, both previously eaten regularly caused immediate symptoms upon reintroduction.

Conclusion

This case demonstrates that in the face of multiple sensitisation to foods, an elemental diet followed by planned and careful staged re-introduction of foods may be effective in establishing which foods can be safely be consumed.

Author details

¹Royal Brompton & Harefield NHS Trust, Rehabilitation & Therapies, London, UK. ²Imperial College, London, Allergy and Clinical Immunology, London, UK.

Published: 12 August 2011

doi:10.1186/2045-7022-1-S1-P90

Cite this article as: Skypala and Till: A case of adult eosinophilic oesophagitis. Clinical and Translational Allergy 2011 1(Suppl 1):P90.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- · No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit



Full list of author information is available at the end of the article



¹Royal Brompton & Harefield NHS Trust, Rehabilitation & Therapies, London,