



ORAL PRESENTATION

Open Access

O19 - Changes in the balance between myeloid (mDC) and plasmacytoid (pDC) cell numbers in peripheral blood during childhood

Paraskevi Maggina^{*}, Vassiliki Papaevangelou, Maria Tsofia, Nikolaos G Papadopoulos

From 3rd Pediatric Allergy and Asthma Meeting (PAAM)
Athens, Greece. 17-19 October 2013

Background

Dendritic cells (DCs) are the most potent antigen-presenting cells, having an important role in linking innate and adaptive immunity. DCs are also critical mediators of immune tolerance and anergy, depending on the type of antigen they encounter. Peripheral blood DCs represent only the 0.1–1% of mononuclear cells and, based on their lineage origins, they can be divided into two major subsets, plasmacytoid DCs (pDCs) and myeloid DCs (mDCs). Although, the total number of all blood circulating lymphocyte subpopulations in children declines with age, previous studies performed on age-related DC changes have shown controversial results [1-4].

Material and methods

Blood samples were obtained from 43 clinically healthy children aged between 1 day to 11 years old during routine examinations for minor elective surgery or routine check-ups in the outpatient clinic. DCs were identified, by FACS-can flow cytometer, as showing no labeling for the “lineage cocktail” (fluorescein isothiocyanate (FITC)-conjugated monoclonal antibodies including CD3, CD14, CD16, CD19, CD20, CD56) and strong labeling for HLA-DR. The percentages of mDCs (CD 11+) and pDCs (CD 123+) were determined using three-colour flow cytometry and their absolute numbers were calculated by using their percentage in relation to the lymphocyte and monocyte number, as determined by differential blood count.

Results

Similarly to previous studies' findings [2,3], we demonstrate that while mDCs do not change with age, pDCs

decrease significantly with age (linear regression: $p=0.0242$, $R^2=0.1343$). Moreover, the mDCs/pDCs ratio showed a significant positive correlation with age during childhood (linear regression: $p=0.0044$, $R^2=0.2092$).

Conclusion

The human immune system is functional less mature during infancy and within the first years of life. Although young children show adult levels of mDCs, the dynamic changes in the balance between mDCs and pDCs during childhood may play a role in the vulnerability of young children to viral and bacterial infections.

Published: 28 February 2014

References

1. Agrawal A, Agrawal S, Tay J, Gupta S: **Biology of dendritic cells in aging.** *J. Clin. Immunol* 2008, **28**:14.
2. Jing Y, Shaheen E, Drake RR, Chen N, Gravenstein S, Deng Y: **Ageing is associated with a numerical and functional decline in plasmacytoid dendritic cells, whereas myeloid dendritic cells are relatively unaltered in human peripheral blood.** *Hum. Immunol* 2009, **70**:777.
3. Teig N, Moses D, Gieseler S, Schauer U: **Age-related changes in human blood dendritic cells subpopulations.** *Scand J Immunol* 2002, **55**:453-7.
4. Heinze A, Elze MC, Kloess S, Ciocarlie O, Königs C, Betz S, Bremm M, Esser R, Klingebiel T, Serban M, Hutton JL, Koehl U: **Age matched dendritic cell subpopulations reference values in childhood.** *Scand J Immunol* 2013, **77**:213-20.

doi:10.1186/2045-7022-4-S1-O18

Cite this article as: Maggina et al.: O19 - Changes in the balance between myeloid (mDC) and plasmacytoid (pDC) cell numbers in peripheral blood during childhood. *Clinical and Translational Allergy* 2014 **4**(Suppl 1):O19.

^{2nd} Department of Pediatrics, “P. and A. Kyriakou” Children’s Hospital, National and Kapodestrian University of Athens, Athens, Greece