



ORAL PRESENTATION

Open Access

O23 - Functional effects of an amino-acid based formula with synbiotics in cow's milk allergic infants

Wesley A Burks¹, Lucien F Harthoorn², Jane E Langford², Marleen T Van Ampting^{2*}, Steven B Goldberg³, Peck Y Ong⁴, Brandon J Essink⁵, Robert B Scott⁶, Bryan M Harvey⁷

From 3rd Pediatric Allergy and Asthma Meeting (PAAM)
Athens, Greece. 17-19 October 2013

Pre- and probiotics (synbiotics) are suggested to have beneficial effects on human health. This study describes effects of an amino acid based formula (AAF) with synbiotics in infants with cow's milk allergy (CMA).

In a prospective, randomized, double-blind controlled study, full term infants with diagnosed IgE and/or non-IgE mediated CMA received a commercially available AAF (NEO; n=56) or an AAF with synbiotics (neutral and acidic oligosaccharides, *Bifidobacterium breve* M-16V) (NEO-SYN; n=54) for 16 weeks. Primary outcome was growth and formula tolerance; secondary outcomes were dermatological/respiratory allergic characteristics, and stool characteristics recorded in subject diaries and evaluated by a physician. Furthermore, adverse events and concomitant medications used during the study were reported.

Average age of infants at inclusion was 4.5±2.4 months. Overall, NEO-SYN and NEO were equally tolerated and both supported normal growth. Both formulas reduced allergic symptoms; SCORAD decreased in both groups with no significant differences in SCORAD change over 16 weeks between both groups. Softer and yellow/brown stools were reported more frequently in the NEO-SYN group compared to more dry and green/dark brown stools in the NEO group (week 0-2, p=0.035; week 2-4, p=0.010 for stool consistency). Furthermore, the NEO-SYN group had less subjects with reported infections (p=0.008) and less subjects receiving medication for functional gastrointestinal disorders (p=0.029) when compared with the NEO group. In addition the NEO-SYN group had a lower

number of infants on antibiotics (p=0.049), especially amoxicillin (p=0.004).

This study shows that an AAF with synbiotics is equally tolerated, supports normal growth and is similarly effective in the management of CMA symptoms compared to an AAF without synbiotics. Infants taking an AAF with this synbiotic blend have different stool characteristics, less reported infections and less antibiotic use, which suggests that addition of the synbiotics to an AAF improves resistance to infections and reduces specific medication use.

Authors' details

¹University of North Carolina, Chapel Hill, NC, USA. ²Nutricia Advanced Medical Nutrition, Danone Research – Centre for Specialised Nutrition, Liverpool, United Kingdom. ³Visions Clinical Research, Tucson, AZ, USA. ⁴Children's Hospital Los Angeles, Los Angeles, CA, USA. ⁵Meridian Clinical Research, Omaha, NE, USA. ⁶Asheboro Research Associates, Asheboro, NC, USA. ⁷Children's Investigational Research Program, LLC (CHIRP), Bentonville, AR, USA.

Published: 28 February 2014

doi:10.1186/2045-7022-4-S1-O23

Cite this article as: Burks et al.: O23 - Functional effects of an amino-acid based formula with synbiotics in cow's milk allergic infants. *Clinical and Translational Allergy* 2014 **4**(Suppl 1):O23.

²Nutricia Advanced Medical Nutrition, Danone Research – Centre for Specialised Nutrition, Liverpool, United Kingdom
Full list of author information is available at the end of the article