

POSTER PRESENTATION

Open Access

# Analysis of the IgE epitope profile of soybean allergen Gly m 4

Felix Husslik<sup>1\*</sup>, Diana Mittag<sup>2</sup>, Christian Seutter von Loetzen<sup>3</sup>, Maximilian Hartl<sup>3</sup>, Lothar Vogel<sup>1</sup>, Barbara Ballmer-Weber<sup>4</sup>, Jörg Kleine-Tebbe<sup>5</sup>, Regina Treudler<sup>6</sup>, Jan-Christoph Simon<sup>6</sup>, Kay-Martin Hanschmann<sup>7</sup>, Andreas Reuter<sup>1</sup>, Michaela Gubesch<sup>1</sup>, Paul Rösch<sup>3</sup>, Stefan Vieths<sup>1</sup>, Thomas Holzhauser<sup>1</sup>, Dirk Schiller<sup>1</sup>

From 5th International Symposium on Molecular Allergology (ISMA 2013)  
Vienna, Austria. 6-7 December 2013

## Background

Individuals with birch pollinosis may show allergic reactions after consumption of soybean-containing food. This is caused by cross-reaction of IgE directed against the major birch pollen allergen Bet v 1 with its homologue Gly m 4 from soybean. Birch pollen allergic subjects without soybean allergy often present IgE binding to Gly m 4.

## Objective

To identify IgE epitopes of Gly m 4 in birch pollen-allergic subjects with and without allergy to soybean.

## Methods

Sera from 33 birch pollen-allergic patients with and without clinical reactivity to soy as determined by oral food challenge or convincing history of soy allergy were included in the study. Specific IgE against Bet v 1 and Gly m 4 was determined by ImmunoCAP. IgE binding of the sera to rGly m 4 and soy extract was tested in western blot analysis. To predict putative IgE epitopes, 20 IgE-binding phage-displayed peptide mimics and heptamers representing the total Gly m 4 amino acid sequence were bioinformatically mapped onto the molecular surface of Gly m 4. To verify predicted epitopes, Gly m 4 variants with substituted epitope residues were expressed in and purified from *E. coli*. Secondary and tertiary structures were assessed by CD and NMR spectroscopy. IgE binding of variants was analyzed by competitive immunoblotting and inhibition ELISA.

## Results

Gly m 4- and Bet v 1-specific IgE levels ranged from 0 to >100 kU<sub>A</sub>/L in sera of both patient groups. CAP values

were above 3.5 kU<sub>A</sub>/L in 18 sera (86%) of patients with clinical soy allergy and in 7 sera (58%) of patients sensitized, but without clinical reactivity to soy, respectively. Intensity of IgE binding in western blot corresponded to ImmunoCAP analysis. Bioinformatic mapping resulted in four distinct epitopes on Gly m 4 surface. Several amino acid residues were chosen for generation of substitutional variants. Purified rGly m 4 variants showed a typical Bet v 1-like structure according to CD spectra and 1D-<sup>1</sup>H-NMR spectroscopy. IgE binding to several substitutional variants was reduced, indicating a multiple number of epitopes on Gly m 4 surface.

## Conclusion

No significant differences in Gly m 4-specific IgE binding could be observed between birch pollen-allergic patients with and without clinical soy allergy in ImmunoCAP and immunoblot analysis. However residual IgE reactivity persists in substitutional rGly m 4 variants, indicating the presence of further IgE epitopes not yet identified.

## Authors' details

<sup>1</sup>Paul-Ehrlich-Institut, Division of Allergology, Langen, Germany. <sup>2</sup>Australian Red Cross Blood Service, Research & Development Division, Melbourne, Australia. <sup>3</sup>University of Bayreuth, Department of Biopolymers, Bayreuth, Germany. <sup>4</sup>University Hospital Zurich, Allergy Unit, Department of Dermatology, Zurich, Switzerland. <sup>5</sup>Allergy and Asthma Center Westend, Praxis Hanf, Ackermann & Kleine-Tebbe, Berlin, Germany. <sup>6</sup>University of Leipzig, Department of Dermatology and Allergology, Leipzig, Germany. <sup>7</sup>Paul-Ehrlich-Institut, Department of Biostatistics, Langen, Germany.

Published: 17 March 2014

doi:10.1186/2045-7022-4-S2-P42

Cite this article as: Husslik et al.: Analysis of the IgE epitope profile of soybean allergen Gly m 4. *Clinical and Translational Allergy* 2014 **4**(Suppl 2):P42.

<sup>1</sup>Paul-Ehrlich-Institut, Division of Allergology, Langen, Germany  
Full list of author information is available at the end of the article