



POSTER PRESENTATION

Open Access

Basophil activation test in allergic patients to platins undergoing rapid desensitization

Pedro Giavina-Bianchi¹, Joana Caiado², Matthieu Picard², Violeta Régnier Galvão³, Mariana Castells^{2*}

From 6th Drug Hypersensitivity Meeting (DHM 6)
Bern, Switzerland. 9-12 April 2014

Background

Desensitization (DST) has become a cornerstone in the management of immediate hypersensitivity reactions (HSRs) to chemotherapeutic agents. It is the only effective procedure for overcoming HSRs to first-line therapy, thus representing an important advance in patients' treatment and prognosis. Nevertheless, there are still no good biomarkers to monitor DST safety and effectiveness. The basophil activation test (BAT) has greater applicability in difficult cases with inconclusive diagnosis and when other diagnostic tests (skin tests and serum specific IgE) have not been standardized, as in drug-induced reactions. The main goal of our study was to assess BAT as a test to monitor rapid DST in patients allergic to platins.

Method

We studied 7 oncologic patients who presented platin allergy and 4 healthy volunteers who had never been exposed to platins. We performed the BAT immediately before DST to platins, assessing CD 203c and CD63 expression on basophils. All patients were evaluated at least in 2 different DST.

Results

BAT was positive in 6 patients (85.7%), with increased expression of CD203c and CD63 in 6 and 3 patients, respectively. Subsequent BAT analysis in different DST procedures showed that the test remained positive with an even greater expression of CD203c and CD63 on basophils after platins exposure. Some patients with positive BAT reacted during DST, in spite of being premedicated, showing the correlation between BAT results and clinical outcomes. Further investigation is necessary to determine BAT predictive values and its clinical applicability.

Conclusion

We developed a BAT to platins that presented good sensitivity in our study. Short-term DST to platinum drugs does not induce persistent hyporesponsiveness on basophils, highlighting the need to maintain DST in allergic patients to platins.

Authors' details

¹Harvard Medical School, Division of Rheumatology, Allergy and Immunology, USA. ²Harvard Medical School, Division of Rheumatology, Allergy and Immunology, Department of Medicine, BWH, USA. ³University of Sao Paulo, Clinical Immunology and Allergy, Brazil.

Published: 18 July 2014

doi:10.1186/2045-7022-4-S3-P34

Cite this article as: Giavina-Bianchi et al.: Basophil activation test in allergic patients to platins undergoing rapid desensitization. *Clinical and Translational Allergy* 2014 **4**(Suppl 3):P34.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



²Harvard Medical School, Division of Rheumatology, Allergy and Immunology, Department of Medicine, BWH, USA
Full list of author information is available at the end of the article